



# Purina® Rangeland® Calf Creep 16 B68

**Specie:** Cattle      **Type of Feed:** Type C Medicated      **Form of Feed:** Pellet

**General Description:** For control of coccidiosis caused Eimeria bovis and E. zuernii in cattle up to 800 lbs. For increased rate of weight gain in pasture cattle (slaughter, stocker, feeder cattle, and dairy and beef replacement heifers.

## PURINA® RANGELAND® CALF CREEP 16 B68 MEDICATED

SUPPLEMENT FEED CALVES FROM BIRTH TO WEANING  
CAUTION: USE ONLY AS DIRECTED

### ACTIVE DRUG INGREDIENTS

Lasalocid ..... 68.00 g/ton

### GUARANTEED ANALYSIS

Crude Protein, (Min) ..... 16.00 %  
 Crude Fat, (Min) ..... 1.00 %  
 Crude Fiber, (Max) ..... 25.00 %  
 Acid Detergent Fiber (ADF), (Max) ..... 12.00 %  
 Calcium (Ca), (Min) ..... 0.50 %  
 Calcium (Ca), (Max) ..... 1.00 %  
 Phosphorus (P), (Min) ..... 0.40 %  
 Salt (NaCl), (Min) ..... 0.25 %  
 Salt (NaCl), (Max) ..... 0.75 %  
 Potassium (K), (Min) ..... 0.70 %  
 Vitamin A, I.U./lb, (Min) ..... 10,000

### INGREDIENTS

Processed Grain By-Products, Roughage Products, Plant Protein Products, Calcium Carbonate, Salt, Vitamin A Supplement, Vitamin E Supplement, Natural and Artificial Flavors, Vitamin D3 Supplement, Cobalt Carbonate, Manganese Sulfate, Ethylenediamine Dihydroiodide, Zinc Sulfate, BHT (a Preservative), Basic Copper Chloride, Sodium Selenite.

### DIRECTIONS

For coccidiosis control hand feed continuously at a rate of 1.34 lbs per 100 lbs of body weight to provide 1 mg lasalocid per 2.2 lbs body weight per day (up to a maximum of 360 mg per head per day).

For increased rate of gain feed continuously at a rate of 1.8 to 8.8 lbs to provide 61 to 300 mg of lasalocid per head per day.

Intakes of lasalocid in excess of 200 mg per head per day have not shown to be more effective than 200 mg per head per day. The drug must be contained in at least 1 pound of feed.

### WARNING

A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

**DO NOT FEED TO SHEEP DUE TO HIGH LEVELS OF SUPPLEMENTAL COPPER.**

Store in a dry, well-ventilated area protected from rodents and insects. Do not feed moldy or insect-infested feed to animals as it may cause illness, performance loss or death.

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed.

USE ONLY AS DIRECTED.

### IMPORTANT:

Follow these management practices:

1. This product is intended only to be fed to suckling calves.
2. This product is not intended to be used as a starter diet or with milk replacer.
3. For maximum intake control this product should be hand fed.
4. If using a creep feeder, make sure the feeder is in good working order.
5. Adjust the slide gate to control feed accumulation at the trough.
6. Troughs should be well protected and managed to prevent the accumulation of fines, wet and moldy feed.
7. Provide adequate feeder space for each animal.
8. Provide mineral and salt free choice.
9. Provide fresh, clean water near the feeding area.