



PRECON[®] COMPLETE

Specie: Beef

Type of Feed: Complete

Form of Feed: Pellet

General Description: Precon[®] Complete Feed with RX3 Immune Support Technology[™] is a complete pelleted ration for preconditioning calves on farm or receiving calves in backgrounding operations or feedlots.

PURINA[®] PRECON[®] COMPLETE

COMPLETE FEED FOR STARTER FEEDLOT CATTLE

GUARANTEED ANALYSIS

Crude Protein, (Min).....	12.00 %
(This includes not more than 2.00% equivalent crude protein from non-protein nitrogen).	
Crude Fat, (Min).....	1.00 %
Crude Fiber, (Max).....	26.00 %
Calcium (Ca), (Min).....	0.45 %
Calcium (Ca), (Max).....	0.95 %
Phosphorus (P), (Min).....	0.40 %
Salt (NaCl), (Min).....	0.25 %
Salt (NaCl), (Max).....	0.75 %
Potassium (K), (Min).....	1.10 %
Vitamin A, I.U./lb, (Min)	5000

INGREDIENTS

Note: ingredients differ by manufacturing plant

Roughage Products, Processed Grain By-Products, Grain Products, Plant Protein Products, Molasses Products, Forage Products, Calcium Carbonate, Lignin Sulfonate, Urea, Salt, Potassium Chloride, Dicalcium Phosphate, Monocalcium Phosphate, Vegetable Oil, Silicon Dioxide, Artificial Flavor, Yeast Extract, Saccharin Sodium, Glycyrrhizin Ammoniated, Sodium Selenite, Sodium Silico Aluminate, Chromium Propionate, Propionic Acid (a Preservative), Propylene Glycol, Zinc Sulfate, Mineral Oil, Dried Enterococcus lactis Fermentation Product, Dried Bacillus licheniformis Fermentation Product, Dried Bacillus subtilis Fermentation Product, Vitamin A Supplement, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper Amino Acid Complex, Cobalt Carbonate, Manganese Sulfate, Cobalt Glucoheptonate, Ethylenediamine Dihydroiodide, Basic Copper Chloride.

DIRECTIONS:

Initially, feed this product at a minimum of 1.5% of body weight and gradually increase the feed rate to 3% of body weight. For example, a 500-pound calf would initially receive a minimum of 7.5 pounds per head per day and gradually increase up to 15 pounds per head per day. Consult your local Purina[®] representative for additional information on how to start cattle on feed.

CAUTION:

DO NOT FEED TO SHEEP DUE TO HIGH LEVELS OF SUPPLEMENTAL COPPER. Store in a dry, well-ventilated area protected from rodents and insects. Do not feed moldy or insect-infested feed to animals as it may cause illness, performance loss or death. USE ONLY AS DIRECTED.

IMPORTANT:

Follow these management practices:

1. Do not allow animals to run out of feed.
2. Feeders/bunks should be well protected and managed to prevent accumulation of fines and wet, moldy feed.
3. When making a ration change, allow 7 to 10 days for animals to adjust to the new ration.
4. Provide adequate feeder space for each animal.
5. Stressed cattle should be placed in clean, uncrowded environments.
6. Provide mineral and salt free choice.
7. Provide fresh, clean water near the feeding area.

In groups of cattle, there are certain animals that experience chronic bloat or other digestive disturbances and consequently, are poor performers. In addition, excess feed consumption, severe weather changes resulting in erratic feed consumption and poorly managed feeding practices can increase the incidence of bloat in all cattle. If bloat occurs, please review feed management practices.