

SteakMaker® BP FORMULATOR R1500

**Species:** Beef**Type of Feed:** Mineral**Form of Feed:** Meal**General Description:**

A high calcium mineral supplement designed for finishing cattle on grain or grain by-product (distiller's grains or gluten feed) diets containing adequate protein.

PURINA®
STEAKMAKER® BP FORMULATOR R1500
MEDICATED

(For Ruminants Only)

SUPPLEMENT FEED FOR GROWER-FINISHER FEEDLOT CATTLE

For improved feed efficiency in cattle fed in confinement for slaughter.

CAUTION: FEED AS DIRECTED

ACTIVE DRUG INGREDIENTS

Monensin (as Monensin sodium)..... 1500.00 g/ton

GUARANTEED ANALYSIS

Calcium (Ca), (Min)..... 28.00 %
Salt (NaCl), (Min)..... 11.00 %
Magnesium (Mg), (Min) 1.00 %
Potassium (K), (Min)..... 0.10 %
Copper (Cu), ppm, (Min).....600
Selenium (Se), ppm, (Min).....12
Zinc (Zn), ppm (Min)2150
Vitamin A, I.U./lb, (Min).....120,000
Vitamin D, I.U./lb, (Min).....30,000
Vitamin E, I.U./lb, (Min).....150
Thiamine, ppm, (Min).....1000

INGREDIENTS

Calcium Carbonate, Salt, Processed Grain By-Products, Magnesium Oxide, Molasses Products, Mineral Oil, Colored with Iron Oxide, Zinc Sulfate, Thiamine Mononitrate, Vitamin E Supplement, Vitamin A Supplement, Cobalt Carbonate, Manganese Sulfate, Ethylenediamine Dihydroiodide, Basic Copper Chloride, Sodium Selenite, Vitamin D3 Supplement, Roughage Products.

FEEDING DIRECTIONS:

Thoroughly mix 40 pounds of this Monensin Type B Medicated feed with 1960 pounds of non-medicated feed to obtain one ton of Type C medicated feed containing 30 g/ton Monensin. Feed the resulting Type C feed continuously at a rate of 20 pounds to provide 300 mg of monensin per head per day.

LIMITATIONS FOR USE:

Feed only to cattle being fed in confinement for slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

CAUTION:

Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of monensin recommended in the feeding directions.

IMPORTANT:

This product is to be fed only to the animal species as directed on this label. Follow these management practices:

1. Do not allow animals to run out of feed.
2. Feeders/bunks should be well protected and managed to prevent accumulation of fines and wet, moldy feed.
3. When making a ration change, allow 7 to 10 days for animals to adjust to the new ration.
4. Provide adequate feeder space for each animal.
5. Provide mineral and salt free choice.
6. Provide fresh, clean water near the feeding area.
7. Have feed before cattle at all times. In groups of cattle, there are certain animals that experience chronic bloat or other digestive disturbances and consequently, are poor performers. In addition, excess feed consumption, severe weather changes resulting in erratic feed consumption, and poorly managed feeding practices can increase the incidence of bloat in all cattle. If bloat occurs, please review feed management practices.

USAGE TIPS:

SteakMaker® BP Formulator is a beef is a high calcium supplement designed to be used in high-energy rations that contain adequate protein. Such rations may contain high grain and/or high grain by-products such as distiller's grains or gluten feed.

Recommended feeding rate of SteakMaker® BP Formulator is 0.3 to 0.4 pounds per head per day. Maximum feeding rate is dependent on the feed additive concentration. All ration adjustments should be made on a gradual basis. SteakMaker® BP Formulator should be thoroughly mixed with non-medicated ration prior to feeding. Follow label directions for mixing instructions.