



SteakMaker® Finisher 50-25 R400

Specie: Beef

Type of Feed: Supplement

Form of Feed: Pellet

General Description:

A 50% protein supplement for finishing cattle on grain and roughage.

PURINA®
STEAKMAKER® FINISHER 50-25 R400

MEDICATED
(For Ruminants Only)

SUPPLEMENT FEED FOR FINISHER FEEDLOT CATTLE
For improved feed efficiency in cattle fed in confinement for slaughter.
CAUTION: FEED AS DIRECTED

ACTIVE DRUG INGREDIENTS

Monensin (as Monensin Sodium) 400.00 g/ton

GUARANTEED ANALYSIS

Crude Protein, (Min).....50.0 %
(This includes not more than 25% equivalent crude protein from non-protein nitrogen).
Crude Fat, (Min).....1.00 %
Crude Fiber, (Max).....15.00 %
Calcium (Ca), (Min).....7.50 %
Phosphorus (P), (Min).....0.50 %
Salt (NaCl), (Min).....4.00 %
Magnesium (Mg), (Min)0.80 %
Potassium (K), (Min).....1.00 %
Copper (Cu), ppm, (Min)200
Selenium (Se), ppm, (Min)4.0
Zinc (Zn), ppm (Min).....700
Vitamin A, I.U./lb, (Min)40,000
Vitamin D, I.U./lb, (Min)10,000
Vitamin E, I.U./lb, (Min).....50

FEEDING DIRECTIONS:

Thoroughly mix 150 pounds of supplement (Type B medicated feed) with 1850 pounds of non-medicated feed to obtain one ton (Type C medicated feed) containing 30 g/ton monensin. Feed the resulting one ton (Type C medicated feed) at a rate of 20 pounds to provide 300 mg monensin per head per day.

LIMITATIONS FOR USE:

Feed only to cattle being fed in confinement for slaughter.

CAUTION:

Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of monensin recommended in the feeding directions.

INGREDIENTS

Note: ingredients differ by manufacturing plant

Processed Grain By-Products, Calcium Carbonate, Urea, Plant Protein Products, Forage Products, Salt, Ammonium Sulfate, Magnesium Oxide, Potassium Chloride, Cobalt Carbonate, Manganese Sulfate, Ethylenediamine Dihydroiodide, Vitamin D3 Supplement, Zinc Sulfate, Basic Copper Chloride, Roughage Products, Sodium Selenite, Vitamin E Supplement, Vitamin A Supplement.

IMPORTANT:

This product is to be fed only to the animal species as directed on this label. Follow these management practices:

1. Provide adequate bunk space for each animal. Feeders/bunks should be well protected and managed to prevent accumulation of fines and wet, moldy feed.
2. When making a ration change, allow 7-10 days for animals to adjust to the new ration.
3. Provide a source of fresh, clean water at all times.
4. Provide mineral and salt free choice.
5. Consult your veterinarian for recommended health program for your local area.
6. Have feed before cattle at all times. In groups of cattle, there are certain animals that experience chronic bloat or other digestive disturbances and consequently, are poor performers. In addition, excess feed consumption, severe weather changes resulting in erratic feed consumption, and poorly managed feeding practices can increase the incidence of bloat in all cattle. If bloat occurs, please review feed management practices.