



SteakMaker® Finisher 50-25

Specie: Beef

Type of Feed: Supplement

Form of Feed: Pellet

General Description:

A 50% protein supplement for finishing cattle on grain and roughage.

PURINA®
STEAKMAKER® FINISHER 50-25
 (For Ruminants Only)
 SUPPLEMENT FEED FOR FINISHER FEEDLOT CATTLE
CAUTION: FEED AS DIRECTED

GUARANTEED ANALYSIS

Crude Protein, (Min).....	50.0 %
(This includes not more than 25% equivalent crude protein from non-protein nitrogen).	
Crude Fat, (Min).....	1.00 %
Crude Fiber, (Max).....	15.00 %
Calcium (Ca), (Min).....	7.50 %
Phosphorus (P), (Min).....	0.50 %
Salt (NaCl), (Min).....	4.00 %
Magnesium (Mg), (Min)	0.80 %
Potassium (K), (Min).....	1.00 %
Copper (Cu), ppm, (Min)	200
Selenium (Se), ppm, (Min)	4.0
Zinc (Zn), ppm (Min).....	700
Vitamin A, I.U./lb, (Min)	40,000
Vitamin D, I.U./lb, (Min)	10,000
Vitamin E, I.U./lb, (Min).....	50

FEEDING DIRECTIONS:

Feed 1 pound per head per day daily along with grain and good quality forage to finishing cattle.

LIMITATIONS FOR USE:

Feed only to cattle being fed in confinement for slaughter.

CAUTION:

Store feed in a dry area away from insects. Do not feed moldy or insect-infested feed to animals as it may cause illness, abortion or death.

INGREDIENTS

Note: ingredients differ by manufacturing plant

Plant Protein Products, Processed Grain By-Products, Calcium Carbonate, Urea, Salt, Molasses Products, Magnesium Oxide, Cobalt Carbonate, Manganese Sulfate, Ethylenediamine, Dihydroiodide, Zinc Sulfate, Vitamin D3 Supplement, Basic Copper Chloride, Roughage Products, Sodium Selenite, Vitamin E Supplement, Vitamin A Supplement.

IMPORTANT:

This product is to be fed only to the animal species as directed on this label. Follow these management practices:

1. Provide adequate bunk space for each animal. Feeders/bunks should be well protected and managed to prevent accumulation of fines and wet, moldy feed.
2. When making a ration change, allow 7-10 days for animals to adjust to the new ration.
3. Provide a source of fresh, clean water at all times.
4. Provide mineral and salt free choice.
5. Consult your veterinarian for recommended health program for your local area.
6. Have feed before cattle at all times. In groups of cattle, there are certain animals that experience chronic bloat or other digestive disturbances and consequently, are poor performers. In addition, excess feed consumption, severe weather changes resulting in erratic feed consumption, and poorly managed feeding practices can increase the incidence of bloat in all cattle. If bloat occurs, please review feed management practices.

USAGE TIPS:

SteakMaker® 50-25 is a beef supplement for finishing cattle. It is ideal for use in moderate to high grain diets. It can fit well in these diets when fed with a wide range of roughage sources. It is typically fed at 0.75 to 1.5 lb. /hd. /d depending on the nutrient profile of other ration ingredients, ration energy content, age and size of cattle being fed and the medicated option used.

All ration adjustments should be made on a gradual basis. Use good bunk management techniques and procedures to prevent digestive disturbances and to optimize dry matter intake and animal performance. SteakMaker® 50-25 should be thoroughly mixed with grain and roughage. Do not top dress or feed separately. Follow label directions for mixing instructions.