



SteakMaker® Grower 36-0 R400

Specie: Beef

Type of Feed: Supplement

Form of Feed: Pellet

General Description:

This pelleted supplement is designed for growing and finishing cattle on moderately high grain diets.

PURINA®

STEAKMAKER® GROWER 36-0 R400

MEDICATED

(For Ruminants Only)

SUPPLEMENT FEED FOR GROWER FEEDLOT CATTLE

For improved feed efficiency in cattle fed in confinement for slaughter.

CAUTION: FEED AS DIRECTED

ACTIVE DRUG INGREDIENTS

Monensin (as Monensin Sodium) 400.00 g/ton

GUARANTEED ANALYSIS

Crude Protein, (Min) 36.00 %
 Crude Fat, (Min) 1.00 %
 Crude Fiber, (Max) 15.00 %
 Calcium (Ca), (Min) 5.00 %
 Phosphorus (P), (Min) 0.80 %
 Salt (NaCl), (Min) 3.50 %
 Magnesium (Mg), (Min) 0.50 %
 Potassium (K), (Min) 0.70 %
 Copper (Cu), ppm, (Min) 135
 Selenium (Se), ppm, (Min) 2.75
 Zinc (Zn), ppm (Min) 467
 Vitamin A, I.U./lb, (Min) 27,000
 Vitamin D, I.U./lb, (Min) 6,700
 Vitamin E, I.U./lb, (Min) 35

INGREDIENTS

Note: ingredients differ by manufacturing plant

Grain Products, Roughage Products (not more than 40%), Calcium Carbonate, Processed Grain By-Products, Salt, Plant Protein Products, Molasses Products, Zinc Sulfate, Manganese Sulfate, Magnesium Oxide, Basic Copper Chloride, Sodium Selenite, Ethylenediamine Dihydriodide, Cobalt Carbonate, Vitamin A Supplement, Vitamin D₃ Supplement, Vitamin E Supplement, Mineral Oil.

FEEDING DIRECTIONS:

Thoroughly mix 150 pounds of this monensin Type B medicated feed into 1850 pounds of non-medicated feed to obtain one ton of Type C medicated feed containing 30 g/ton monensin.

FEEDING DIRECTIONS:

Feed the resulting Type C medicated feed continuously at a rate of 16.7 pounds to provide 250 mg of monensin per head per day.

WARNING:

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

CAUTION:

Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of monensin recommended in the feeding directions.

IMPORTANT:

This product is to be fed only to the animal species as directed on this label. Follow these management practices:

1. Do not allow animals to run out of feed.
2. Provide adequate bunk space for each animal. Feeders/bunks should be well protected and managed to prevent accumulation of fines and wet, moldy feed.
3. When making a ration change, allow 7-10 days for animals to adjust to the new ration.
4. Provide a source of fresh, clean water at all times.
5. Provide mineral and salt free choice.
6. In groups of cattle, there are certain animals that experience chronic bloat or other digestive disturbances and consequently, are poor performers. In addition, excess feed consumption, severe weather changes resulting in erratic feed consumption, and poorly managed feeding practices can increase the incidence of bloat in all cattle. If bloat occurs, please review feed management practices.

USAGE TIPS:

SteakMaker® Grower supplement is pelleted, moderate calcium designed to be used in growing/finishing rations.

Recommended feeding rates of SteakMaker® Grower supplement is 1.5 pounds per head per day. Thoroughly mix SteakMaker® Grower supplement with the non-medicated ration prior to feeding.