



PURINA® SUPERMIX B1800

Specie: Cattle and Sheep **Type of Feed:** Supplement **Form of Feed:** Pellet

PURINA® SUPERMIX B1800

TYPE B MEDICATED FEED
SUPPLEMENT FEED FOR CATTLE AND SHEEP

For improved feed efficiency in cattle fed in confinement for slaughter. For improved feed efficiency and increased rate of weight gain in cattle fed in confinement for slaughter. For increased rate of weight gain in pasture cattle (slaughter, stocker, feeder cattle and dairy and beef replacement heifers). For control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in cattle up to 800 lbs. For the prevention of coccidiosis caused by *Eimeria ovina*, *E. crandallis*, *E. Ovinoidalis* (*E. ninakohlyakimovae*), *E. Parva* and *E. intricata* in sheep maintained in confinement.

ACTIVE DRUG INGREDIENTS

Lasalocid (as Lasalocid sodium) 1800.00 g/ton

GUARANTEED ANALYSIS

Crude Protein, (Min) 8.00 %

Crude Fat, (Min) 1.00 %

Crude Fiber, (Max) 30.00 %

INGREDIENTS

Note: ingredients differ by manufacturing plant

Roughage Products, Processed Grain By-Products, Molasses Products.

DIRECTIONS:

Each pound of this product contains 0.9 g (90 mg) lasalocid. Use the following chart to mix this Type B medicated feed with non-medicated feed to obtain one ton of Type C Medicated feed.

Mixing Chart:		
Pounds of Supermix B1800 (lbs)	Pounds of non-medicated feed (lbs)	Resulting Type C Feed lasalocid level (g/ton)
11.1	1988.9	10
22.2	1977.8	20
27.8	1972.2	25
33.3	1966.7	30
66.7	1933.3	60
100.0	1900.0	90
222.2	1777.8	200

For improved feed efficiency in cattle fed in confinement for slaughter: Use the above mixing chart to create a finished feed containing 10 to 30 g/ton lasalocid. Feed mixed feed continuously to provide not less than 100 mg or more than 360 mg lasalocid/head/day.

DIRECTIONS (CONTINUED):

For improved feed efficiency and increased rate of weight gain in cattle fed in confinement for slaughter: Use the above mixing chart to create a finished feed containing 25 to 30 g/ton lasalocid. Feed mixed feed continuously to provide not less than 250 mg or more than 360 mg lasalocid/head/day.

For increased rate of weight gain in pasture cattle (slaughter, stocker, feeder cattle and dairy and beef replacement heifers): Use the above mixing chart to create a finished feed containing 60 to 200 g/ton lasalocid. Feed mixed feed continuously at a rate of not less than 60 mg or more than 300 mg of lasalocid/head/day. Intakes of lasalocid in excess of 200 mg per head per day have not been shown to be more effective than 200 mg per head per day. This product must be mixed with grain and/or forage and fed in at least 1 lb of feed.

For control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in cattle weighing up to 800 lbs: Use the above mixing chart to create a finished feed containing 90 g/ton lasalocid. Hand feed the mixed feed continuously to provide 1 mg lasalocid per 2.2 lbs body weight per day (up to a maximum of 360 mg per head per day). 1 lb of feed will medicate 100 lbs of body weight.

For prevention of coccidiosis caused by *Eimeria ovina*, *E. crandallis*, *E. ovinoidalis* (*E. ninakohlyakimovae*), *E. parva* and *E. intricate* in sheep maintained in confinement: Use the above mixing chart to create a finished feed containing 20 to 30 g/ton lasalocid of complete feed. Feed the complete feed to provide no less than 15 mg or more than 70 mg lasalocid/head/day depending on body weight.

CAUTION:

Feeding undiluted or mixing errors may result in an excess lasalocid concentration which could be fatal to cattle and sheep. Do not allow horses or other equines access to Type A articles or Type B feeds containing lasalocid as ingestion may be fatal. Safety of lasalocid for use in unapproved species has not been established. Store in a dry, well-ventilated area protected from rodents and insects. Do not feed moldy or insect-infested feed to animals as it may cause illness, performance loss or death. **USE ONLY AS DIRECTED**